



Public Health
Prevent. Promote. Protect.

Kansas City, MO, Health Dept.

Osteoporosis

Osteoporosis is a common and costly problem that could become even more widespread. According to *The 2004 Surgeon General's Report on Bone Health and Osteoporosis*, by 2020, half of all men and women ≥ 50 y of age will have osteoporosis if they do not change their diet and lifestyle.³⁰⁵ Some 10 million individuals are estimated to already have the disease and almost 34 million more are estimated to have low bone mass (osteopenia), placing them at risk of osteoporosis. Women are four times more likely than men to develop the disease, 8 million women compared to 2 million men. In a 2006 telephone survey commissioned by the Kansas City Health Department 10% of 1,229 households had at least one person afflicted with osteoporosis and that 82% of these individuals were female.³⁰⁶

Osteoporosis is a disease in which bones become fragile and more likely to break. If not prevented or if left untreated, osteoporosis can progress painlessly until a bone breaks. The disease is responsible for >1.5 million fractures annually and these occur typically in the hip (>300,000), spine (700,000), and wrist (250,000); ~300,000 fractures occur in other bones. The National Osteoporosis Foundation (www.nof.org) estimates that by 2010, more than 1 million Missourians will have osteoporosis or osteopenia.

Any bone can be affected, but of special concern are fractures of the hip and spine. A hip fracture almost always requires hospitalization and major surgery. It can impair a person's ability to walk unassisted and may cause prolonged or permanent disability or even death. Spinal or vertebral fractures also have serious consequences, including loss of height, severe back pain, and deformity. Each year in the US osteoporotic fractures lead to >500,000 hospitalizations, >800,000 emergency department encounters, >2,600,000 physician office visits, and the placement of nearly 180,000 individuals into nursing homes.

Nationally, in 2002, direct expenditures by hospitals and nursing homes for osteoporotic hip fractures

³⁰⁵ US Department of Health and Human Services. 2004. *Bone Health and Osteoporosis: A Report of the Surgeon General*. Atlanta, GA. Department of HHS, CDC, NCCDPHP, Office of the Surgeon General.

³⁰⁶ Kansas City Health Department. 2006 Health Planning and Assessment Survey. www.kcmo.org.



were \$18 billion. The Missouri Department of Health and Senior Services reported that, in 2003, approximately 7,200 Missourians were hospitalized due to a hip fracture and that hospital costs related to those fractures exceeded \$180 million.

An analysis conducted by the Kansas City Health Department found that in 2003 there were 1,238 emergency department visits and 1,139 hospitalizations for osteoporotic fractures among City residents ≥ 50 y of age. Hip fractures accounted for 43% of the hospitalizations, with 79.5% of the hip fractures being among persons ≥ 75 y old. Overall, direct costs for osteoporotic fractures totaled \$1,680,612 for emergency department visits and \$31,022,374 for hospitalizations. Hip fractures accounted for \$16,114,090 or 52% of the hospitalization costs.

Significant risk has been reported in people of all ethnic backgrounds. A white woman over the age of 50 has a $>40\%$ chance of suffering a fracture sometime during the rest of her life. While the lifetime risk for men and non-white women is less across all types of fractures, it is nonetheless substantial and may be rising in some groups such as Hispanic women.

Twenty percent of non-Hispanic white women and Asian women ≥ 50 years of age are estimated to have osteoporosis and 52% are estimated to have osteopenia. For Hispanic women the estimates are 10% and 49%, respectively, while for non-Hispanic black women the estimates are 5% and 35%, respectively. It is believed that that osteoporosis is both under-recognized and under-treated in both white and black women.

Among men the estimates for both osteoporosis and low bone mass are lower than for women,³⁰⁷ although the incidence and costs of fractures in men is rising (currently estimated at 30% of total costs for treating fragility fractures). For non-Hispanic white and Asian men ≥ 50 years of age, an estimated 7% have osteoporosis and 35% have osteopenia ≥ 45 . For Hispanic men the estimates are 3% and 23% respectively, while for non-Hispanic black men they are 4% and 19%, respectively.

Fractures can have devastating consequences for both the individuals who suffer them and family members. For example, the Surgeon General's report points out that hip fractures are associated with

³⁰⁷ Melton LJ et al. 1998. Bone density and fracture risk in men. *J Bone Mineral Res* 13:1915-1923.

OSTEOPOROSIS



Public Health
Prevent. Promote. Protect.

Kansas City, MO, Health Dept.

increased risk of mortality. The risk of mortality is 2.8-4 times greater among hip fracture patients during the first 3 months after the fracture, as compared to the comparable risk among individuals of similar age who live in the community and do not suffer a fracture. Those persons in poor health or living in a nursing home at the time of fracture are particularly vulnerable. For those that do survive, these fractures often precipitate a downward spiral in physical and mental health that dramatically impairs quality of life. Nearly 20% of hip fracture patients end up in a nursing home.

Osteoporosis is not a natural part of the aging process and can be prevented or detected early and effectively treated. Left unchecked, the bone health status of Americans is only going to get worse, due primarily to the aging of the population. Therefore, a major message of the Surgeon General's report was that the bone health status of Americans can be improved, but much of what could be done to reduce this burden is not being done today. Physical activity and adequate calcium and vitamin D intake are known to be major contributors to bone health for individuals of all ages. Even though bone disease often strikes late in life, the importance of beginning prevention at a very young age and continuing it throughout life is well understood. Improvements in assessment tools have made it possible to detect bone disease early and to identify those at highest risk of fracture. And, therapeutic advances in bone disease have equaled if not surpassed advances in the areas of prevention and diagnosis. The tremendous potential offered by these developments in bone health, unfortunately, has yet to become a reality.